

Radiation from matter entrainment in astrophysical jets: the AGN case

A.T. Araudo^{1,2}, V. Bosch-Ramon³ and G.E. Romero^{1,2}

¹ Instituto Argentino de Radioastronomía (CCT La Plata, CONICET),
 C.C.5, 1894 Villa Elisa, Buenos Aires, Argentina,
 email: aaraudo@fcaglp.unlp.edu.ar, romero@fcaglp.unlp.edu.ar

² Facultad de Ciencias Astronómicas y Geofísicas, Universidad Nacional de La Plata,
 Paseo del Bosque, 1900 La Plata, Argentina

³ Dublin Institute for Advanced Studies, 31 Fitzwilliam Place, Dublin 2, Ireland
 email: valenti@cp.dias.ie

Abstract. Jets are found in a variety of astrophysical sources, from young stellar objects to active galactic nuclei. In all the cases the jet propagates with a supersonic velocity through the external medium, which can be inhomogeneous, and inhomogeneities could penetrate into the jet. The interaction of the jet material with an obstacle produces a bow shock in the jet in which particles can be accelerated up to relativistic energies and emit high-energy photons. In this work, we explore the active galactic nuclei scenario, focusing on the dynamical and radiative consequences of the interaction at different jet heights. We find that the produced high-energy emission could be detectable by the current γ -ray telescopes. In general, the jet-clump interactions are a possible mechanism to produce (steady or flaring) high-energy emission in many astrophysical sources in which jets are present.

Keywords. galaxies: active, radiation mechanisms: nonthermal, gamma rays: theory

1. Introduction

Jets at different scales are present in astrophysical sources such as protostars, microquasars (MQ) and active galactic nuclei (AGN). The medium that surrounds the jets is not homogeneous and clumps from the external medium can interact with them. In young stellar objects (YSO), the interaction of matter from the external medium with the outflows that emanates from the protostar has been proposed to explain the formation of Herbig-Haro (HH) objects. The dynamical properties of the interaction has been simulated by Raga *et al.* (2003). In high mass microquasars (HMMQ), the interaction of clumps from the companion stellar wind (Owocki & Cohen (2006)) with the jets of the compact object can produce significant amount of γ rays in the form of flaring and steady emission (Araudo *et al.* (2009)). In the case where only few clumps interact with the jet, the produced emission is sporadic and can explain the GeV flares detected from some γ -ray binaries (e.g. Abdo *et al.* (2009)).

At extragalactic scales, active galaxies are composed by an accreting supermassive black hole (SMBH) at the center and powerful jets (Begelman *et al.* (1984)). Surrounding the SMBH there are a population of clouds (Krolik *et al.* (1981)) that moves at velocities $> 1000 \text{ km s}^{-1}$ forming the called broad line region (BLR). In the present contribution, we illustrate the most important dynamical and radiative consequences of the interaction of clouds from the BLR with the base of AGN jets.

2. The jet-cloud interaction

We adopt clouds with density $n_c = 10^{10} \text{ cm}^{-3}$, size $R_c = 10^{13} \text{ cm}$, and velocity $v_c = 10^9 \text{ cm s}^{-1}$. The jet Lorentz factor is fixed to $\Gamma = 10$, implying $v_j \approx c$, and the radius/height relation is fixed to $R_j = 0.1 z$. The jet density n_j in the laboratory RF can be estimated as $n_j = L_j / ((\Gamma - 1) m_p c^3 \sigma_j)$, where $\sigma_j = \pi R_j^2$ and L_j is the kinetic power of the matter-dominated jet.

For a completely penetration of a cloud into the jet, the ram pressure of the later should not destroy the former before the cloud has fully entered into the jet. This means that the time required by the cloud to penetrate into the jet, $t_c \sim 2R_c/v_c = 2 \times 10^4 \text{ s}$, should be shorter than the cloud lifetime inside the jet. To estimate this cloud lifetime, we first compute the time required by the shock in the cloud to cross it (t_{cs}). The velocity of this shock is $v_{cs} \sim \chi^{-1/2} c$, where $\chi = n_c/n_j(\Gamma - 1)$, is derived by ensuring that the jet and the cloud shock ram pressures are equal. This yields a cloud shocking time of

$$t_{cs} \sim \frac{2R_c}{v_{cs}} \simeq 7 \times 10^4 \left(\frac{R_c}{10^{13} \text{ cm}} \right) \left(\frac{n_c}{10^{10} \text{ cm}^{-3}} \right)^{1/2} \left(\frac{z}{10^{16} \text{ cm}} \right) \left(\frac{L_j}{10^{44} \text{ erg s}^{-1}} \right)^{-1/2} \text{ s}. \quad (2.1)$$

Rayleigh-Taylor (RT) and Kelvin-Helmholtz (KH) instabilities produced by the interaction with the jet material will affect the cloud. The timescale in which these instabilities grows up to a scale length $\sim R_c$ is $\sim t_{cs}$. For this reason, we take t_{cs} as the characteristic timescale of our study. Therefore, for a penetration time (t_c) at least as short as $\sim t_{cs}$, the cloud will remain an effective obstacle for the jet flow. Setting $t_c \sim t_{cs}$, we obtain the minimum value for the interaction height z_{int} , giving

$$z_{\text{int}}^{\text{min}} \approx 2.5 \times 10^{15} \left(\frac{v_c}{10^9 \text{ cm s}^{-1}} \right)^{-1} \left(\frac{n_c}{10^{10} \text{ cm}^{-3}} \right)^{-1/2} \left(\frac{L_j}{10^{44} \text{ erg s}^{-1}} \right)^{1/2} \text{ cm}. \quad (2.2)$$

For $z_{\text{int}} > z_{\text{int}}^{\text{min}}$, the jet crossing time t_j , characterized by $t_j \sim 2R_j/v_c \sim 5 \times 10^5 \text{ s}$, is larger than t_{cs} . Once the cloud is inside the jet, a bow shock in the later is formed on a time $t_{bs} \sim Z/v_j \sim 10^2 \text{ s}$ at a distance $Z \sim 0.3R_c \text{ cm}$ from the cloud. In this bow shock, particles can be accelerated up to relativistic energies more efficiently than in the cloud shock ($v_{bs} \gg v_{cs}$).

3. Non-thermal emission

We focus here on the particle acceleration in the bow shock. The non-thermal luminosity of particles accelerated in the bow shock can be estimated as a fraction η of the bow shock luminosity: $L_{\text{nt}} \sim \eta_{\text{nt}} L_{\text{bs}}$, where $L_{\text{bs}} \sim (\sigma_c/\sigma_j) L_j$ and $\sigma_c = \pi R_c^2$. The accelerator/emitter magnetic field in the bow-shock RF (B) can be determined by relating $U_B = \eta_B U_{\text{nt}}$, where $U_B = B^2/8\pi$ and $U_{\text{nt}} = L_{\text{nt}}/(\sigma_c c)$ are the magnetic and the non-thermal energy densities, respectively. Fixing $\eta_{\text{nt}} = 0.1$ and $\eta_B = 0.01$, we obtain $L_{\text{nt}} \sim 4 \times 10^{39} (L_j/10^{44} \text{ erg s}^{-1}) \text{ erg s}^{-1}$ and $B \sim 10 \text{ G}$. Regarding the acceleration mechanism, we adopt the following prescription for the acceleration rate: $\dot{E}_{\text{acc}} = 0.1 q B c$.

We assume that electrons and protons are injected into the bow shock region following a power law in energy of index 2.2 with an exponential cutoff at the maximum electron energy. The injection luminosity is L_{nt} . Particles are affected by different losses that balance the energy gain from acceleration. The escape time downstream from the relativistic bow shock consider advection ($t_{\text{adv}} \sim 3 R_c/c \sim 10^3 \text{ s}$) and diffusion ($t_{\text{diff}} = 3 Z^2/2 r_g c$) timescales. In the later, r_g is the particle gyroradius and the Bohm regime has been considered.




Figure 1. Left: acceleration gain, escape, and cooling lepton timescales are plotted. SSC is plotted when the steady state is reached and EC for the both BLR and disc photon fields are shown for the conditions of faint (BLR: 10^{44} ; disc: 10^{45} erg s $^{-1}$) and bright sources (BLR: 10^{46} ; disc: 10^{47} erg s $^{-1}$). Synchrotron and relativistic bremsstrahlung are also plotted. Right: Upper limits to the γ -ray luminosity produced by N_c^j clouds inside the jet as a function of L_j in FR II sources. Two cases are plotted, one assuming that clouds cross the jet without disruption (green solid lines), and one in which the clouds are destroyed in a time as short as t_{cs} (green dashed lines). In addition, the sensitivity levels of *Fermi* in the range 0.1–1 GeV (maroon dotted lines) are plotted for three different distances $d = 10, 100$, and 1000 Mpc.

The most important radiative losses that affect the lepton injection are synchrotron and SSC, determining a maximum energy $E_e^{\max}$ of several TeV, as is shown in Figure 3 (left). In the case of protons, they will not cool efficiently in the bow shock via pp interactions and the maximum energy is constrained by equating the acceleration and diffusion timescales, given $E_p^{\max} \sim 5 \times 10^3 (B/10 \text{ G})(R_c/10^{13} \text{ cm})$ TeV. Then, protons with energies $> 0.4 E_p^{\max}$ can reach the cloud by diffusion and radiate via pp more efficiently than in the jet ($n_c \gg n_j(z_{\text{int}})$).

4. Many clouds interacting with the jets

Clouds fill the BLR, and many of them can be simultaneously inside the jet at different z , each of them producing non-thermal radiation. Therefore, the total luminosity can be much larger than that produced by just one interaction, which is $\sim L_{\text{nt}}$. The number of clouds within the jets, at $z_{\text{int}}^{\min} \leq z \leq R_{\text{blr}}$, can be computed from the jet (V_j) and cloud (V_c) volumes, resulting in

$$N_c^j = 2f \frac{V_j}{V_c} \sim 9 \left(\frac{L_j}{10^{44} \text{ erg s}^{-1}} \right)^2 \left(\frac{R_c}{10^{13} \text{ cm}} \right)^{-3}, \quad (4.1)$$

where the factor 2 accounts for the two jets and $f \sim 10^{-6}$ is the filling factor of clouds in the whole BLR (Dietrich et al. 1999). In reality, N_c^j is correct if one neglects that the cloud disrupts and fragments, and eventually dilutes inside the jet. For instance, Klein et al. (1994) estimated a shocked cloud lifetime of several t_{cs} , and Shin et al. (2008) found that even a weak magnetic field in the cloud can significantly increase its lifetime. Finally, even under cloud fragmentation, strong bow shocks can form around the cloud fragments before these have accelerated close to v_j . All this makes the real number of interacting clouds inside the jet difficult to estimate, but it should be between $(t_{cs}/t_j) N_c^j$ and N_c^j .

The presence of many clouds inside the jet implies that the total non-thermal lumi-

nosity available in the BLR-jet intersection region is

$$L_{\text{nt}}^{\text{tot}} \sim 2 \int_{z_{\text{int}}^{\text{min}}}^{R_{\text{blr}}} \frac{dN_{\text{c}}^{\text{j}}}{dz} L_{\text{nt}}(z) dz \sim 2 \times 10^{40} \left(\frac{\eta_{\text{nt}}}{0.1} \right) \left(\frac{R_{\text{c}}}{10^{13} \text{ cm}} \right)^{-1} \left(\frac{L_{\text{j}}}{10^{44} \text{ erg s}^{-1}} \right)^{1.7}, \quad (4.2)$$

where $dN_{\text{c}}^{\text{j}}/dz$ is the number of clouds located in a jet volume $dV_{\text{j}} = \pi (0.1z)^2 dz$. In both Eqs. (4.1) and (4.2), L_{blr} has been fixed to $0.1 L_{\text{j}}$, as approximately found in FR II galaxies, and R_{blr} has been derived from Kaspi *et al.* (2005).

In Fig. 3 (right), we show estimates of the γ -ray luminosity when many clouds interact simultaneously with the jet. For this, we have followed a simple approach assuming that most of the non-thermal luminosity is converted into γ rays. This will be the case as long as the escape and synchrotron cooling time are longer than the IC cooling time (EC+SSC) at the highest electron energies. Given the little information about the BLR available for FR I sources, we do not specifically consider these sources here. In Araudo *et al.* (2010), we present more detailed calculations by applying the model presented in the present contribution to two characteristic sources, Cen A (FR I, one interaction) and 3C 273 (FR II, many interactions).

5. Discussion

We have studied the interaction of clouds from the BLR with the base of jets in AGNs. For very nearby sources, such as Cen A, the interaction of large clouds with jets may be detectable as a flaring event, although the number of these large clouds and thereby the duty cycle of the flares are difficult to estimate. Given the weak external photon fields in these sources, VHE photons can escape without experiencing significant absorption. Therefore, jet-cloud interactions in nearby FR I may be detectable in both the HE and the VHE range as flares with timescales of about one day.

In FR II sources, many BLR clouds could interact simultaneously with the jet. The number of clouds depends strongly on the cloud lifetime inside the jet, which could be of the order of several t_{cs} . Nevertheless, we note that after cloud fragmentation many bow shocks may still form and efficiently accelerate particles if these fragments move more slowly than the jet. Since FR II sources are expected to exhibit high accretion rates, radiation above 1 GeV produced in the jet base can be strongly attenuated by the dense disc and the BLR photon fields, although γ rays below 1 GeV should not be affected significantly. Since jet-cloud emission should be rather isotropic, it would be masked by jet beamed emission in blazar sources, although since powerful/nearby FR II jets do not display significant beaming, these objects may emit γ rays from jet-cloud interactions. As shown in Fig. 3 (right), close and powerful sources could be detectable by deep enough observations of *Fermi*.

References

- Abdo, A.A. et al. 2009 [arXiv:0902.1340]
- Araudo, A.T., Bosch-Ramon, V., & Romero, G.E. 2009, *A&A*, 503, 673
- Araudo, A.T., Bosch-Ramon, V., & Romero, G.E. 2010, *A&A* (in press) [arXiv:1007.2199]
- Begelman, M.C., Blandford, R.D., & Rees, M.J. 1984, *Rev. Mod. Phys.*, 56, 255
- Kaspi, S., Maoz, D., Netzer, H., Peterson, B.M., Vestergaard, M., & Jannuzi, B.T. 2005, *ApJ*, 629, 61
- Krolik, J.H., McKee, C.F., & Tarter, C.B. 1981, *ApJ*, 249, 422
- Owocki, S.P., & Cohen D.H. 2006, *ApJ*, 648, 5650
- Raga, A.C., Velazquez, P.F., de Gouveia dal Pino, E.M., Noriega-Crespo, A., & Mininni, P. 2003, *RMxAC*, 15, 115